arXiv:1211.4274v1 [math.CA] 18 Nov 2012

INVERSE SPECTRAL THEOREM AND POINT PERTURBATIONS OF EVENTUALLY PERIODIC JACOBI MATRICES

ROSTYSLAV KOZHAN

Abstract. We discuss properties of resonances for exponentially small perturbations of periodic and free Jacobi matrices. We show that eventually periodic Jacobi matrices are uniquely determined by the locations of eigenvalues and resonances, and we fully characterize all possible configurations of eigenvalues and resonances.

We show that it is always possible to add/remove a point mass to/from the spectrum of a Jacobi matrix without decreasing the exponential rate of proximity from the isospectral torus. This generalizes the result of Geronimo [3].

1. Introduction

Periodic Jacobi matrices are the operators on $\ell_2(\mathbb{Z}_+)$ of the form

$$\mathcal{J} = \begin{pmatrix} b_1 & a_1 & 0 & \\ a_1 & b_2 & a_2 & \ddots \\ 0 & a_2 & b_3 & \ddots \\ & \ddots & \ddots & \ddots \end{pmatrix} \tag{1.1}$$

for which there exists some (period) $p \ge 1$ such that

$$a_{n+p} = a_n, \quad b_{n+p} = b_n \quad \text{for all } n. \tag{1.2}$$

Any matrix of the form (1.1) will also be denoted by $(a_n, b_n)_{n=1}^\infty$. Sequences $\{a_n\}$, $\{b_n\}$ of real numbers are called Jacobi parameters.

We will be concerned here with two classes of perturbations of periodic Jacobi matrices. The first contains the eventually periodic matrices, that is, all matrices $\mathcal{J}$ for which there exists $s \ge 0$ such that

$$a_n = a_n^{(0)}, \quad b_n = b_n^{(0)} \quad \text{for all } n \ge s+1, \tag{1.3}$$

for some periodic Jacobi matrix $(a_n^{(0)}, b_n^{(0)})_{n=1}^\infty$.

The second class consists of the exponentially small perturbations of periodic matrices, i.e., those matrices for which there exists $1 < R \le \infty$ such that

$$\limsup_{n\to\infty} \left(|a_n - a_n^{(0)}| + |b_n - b_n^{(0)}| \right)^{1/2n} \le R^{-1}, \tag{1.4}$$

Date: November 20, 2012.

E-mail: kozhan@math.ucla.edu

Address: University of California, Los Angeles; Department of Mathematics; Los Angeles, CA 90095, USA.

for some periodic Jacobi matrix $(a_n^{(0)}, b_n^{(0)})_{n=1}^\infty$.

Let μ be the spectral measure of $\mathcal{J}$ with respect to the vector $(1, 0, 0, \dots)^T$ (we are assuming $a_j > 0$; therefore this vector is cyclic).

It is well-known that essential spectrum of a p-periodic Jacobi matrix consists of p closed intervals which are allowed to touch but otherwise are disjoint. Thus

$$\sigma_{ess}(\mathcal{J}) = \operatorname{ess\,supp} \mu = \bigcup_{j=1}^{l+1} [\alpha_j, \beta_j] \equiv \mathfrak{e}, \quad \alpha_j < \beta_j < \alpha_{j+1}. \tag{1.5}$$

We will refer to $[\alpha_j, \beta_j]$ as a "band", and to (β_j, α_{j+1}) as a "gap". l here is the number of gaps (in general $l \le p - 1$). We call $\mathfrak{e}$ a finite gap set.

Note that the perturbations (1.3), (1.4) that we are considering are compact. Thus their spectrum also satisfies (1.5).

The important special case of periodic Jacobi matrices is with $p = 1$ ($l = 0$). Then the spectrum consists of one band. Typically one normalizes, $a_n = 1$, $b_n = 0$ for all n, in which case the spectrum is $[-2, 2]$. We will refer to this matrix as the "free Jacobi matrix", and its finite rank perturbations will be called "eventually free".

To any μ we can associate

$$m(z) = \int \frac{d\mu(x)}{x - z}, \quad z \notin \operatorname{ess\,supp} \mu, \tag{1.6}$$

the Borel/Stieltjes transform of μ. We will refer to this function as the m-function of $\mathcal{J}$. This is a meromorphic function on $\mathbb{C} \setminus \mathfrak{e}$. As was shown in [10] (see Lemmas 2.5 and 2.6 below), under (1.3) or (1.4), the m-function has a meromorphic continuation through the bands of $\mathfrak{e}$ to some domain of the second sheet of the natural Riemann surface $\mathcal{S}_\mathfrak{e}$ (see Definition 2.1).

In accordance with the continuous Schrōdinger operators, we define resonances to be the poles of m on the second sheet of $\mathcal{S}$. Resonances that are real will be called antibound states. In Section 3 we establish properties of eigenvalues and resonances. In particular we show that between each two consecutive eigenvalues in the same gap there must be an odd number of antibound states. This property has been established by Simon in [13] for Schrōdinger operator with exponentially decaying potentials (analogue of (1.4)). In fact, we show that a stronger fact holds true: if counted from any band edge, every second singularity must be a resonance. See Theorem 3.1 for the rigorous statement.

In Section 4 we solve the inverse problem: the location of eigenvalues and resonances uniquely determines the eventually periodic Jacobi matrix. Moreover, we find the necessary and sufficient conditions on the system of eigenvalues + resonances that can occur for eventually periodic Jacobi matrices. In particular this allows to construct an (eventually periodic) Jacobi matrix with spectrum $\mathfrak{e} \cup \{z_j\}_{j=1}^N$ for any finite sequence $\{z_j\}_{j=1}^N$ of points outside of $\mathfrak{e}$. This result seems to be new even for $l = 0$ case.

In [1] Damanik and Simon found a necessary and sufficient condition for an analytic function u to be the Jost function of a Jacobi matrix with exponentially converging parameters (i.e., in our notation, exponentially small perturbations of $l = 0$ case). These conditions (see Lemmas 5.2 and 5.3) involve a not-so-intuitive notion of canonicity. Using our observations from Sections 3 and 4, we can restate the notion of canonicity entirely in terms of configuration of the zeros of u. This is done in Section 5.

In Section 6 we study the effect that removal or addition of a point mass to the measure (in the literature this is sometimes called the Uvarov transform) has on the Jacobi parameters. We show that in the $l = 0$ case the effect is always exponentially small, and we can calculate the exponential rate of the change (this is not true in general though, see Wong's [16]). When $l > 0$, the effect is no longer exponentially small. However, if $\mathcal{J}$ is exponentially close to a periodic Jacobi matrix $\mathcal{J}_0$, then after the Uvarov transform the new Jacobi matrix will be exponentially close to another periodic Jacobi matrix $\widehat{\mathcal{J}}_0$ with $\sigma_{ess}(\widehat{\mathcal{J}}_0) = \sigma_{ess}(\mathcal{J}_0)$. One may want to think that the effect is still exponentially small, but with respect to the distance from the isospectral torus (see Definition 2.3).

Using our methods, studying the effect of the Christoffel transforms is also easy (by Christoffel transform we will mean the "multiplication" of the measure by a rational function). We do this in Section 7. In particular we show that using a Christoffel transform and a sequence of Uvarov transforms we can always add any finite set of point masses to the spectrum of the measure without decreasing the rate of exponential proximity (in the sense of (1.4)) from the isospectral torus. In fact we can do this for the resonances too, but only if it does not violate the "interlacing" properties of Theorem 3.1 of course. This completely solves the problem studied by Geronimo ($l = 0$) in [3] (see also Geronimo–Nevai [4]): we remove the positivity constraint in [3] that was implicitly imposed on the expression for the weight (see Remark 3 after our Theorem 7.1 or [3, Eq. (3.14)]).

Finally in Section 8 we show that some of the results from Sections 6 and 7 can be generalized to the matrix-valued setting (i.e., when a_n, b_n are square $m \times m$ matrices).

During the write-up of the paper the author stumbled upon the recent paper by Iantchenko–Korotyaev [7] who also study eventually periodic Jacobi matrices using entirely different methods. The only result that seem to be common is a part of our Theorem 3.2 (indeed, the function F from [7] is our function a from Section 4, up to a multiplicative constant). For the inverse problem they use the Jost function rather than m-function, and it does not seem to be straightforward to derive one result from another. Moreover, their methods do not seem to indicate that resonances and eigenvalues uniquely recover the Jacobi matrix. The inverse problem of [7] is also different from ours: they consider finite rank perturbations of a *fixed* periodic Jacobi matrix, while our approach is to classify the spectral data for the eventually periodic Jacobi matrices with a given essential spectrum, i.e., finite rank perturbations of a matrix from the *given isospectral torus*.

Other than the paper [7] (and [1, 12] for $l = 0$), the author is not familiar with any other paper which studies the spectral problem (direct or inverse) of eventually periodic Jacobi matrices. There are many results that treat resonances of the Schrōdinger operator, as well as the full line Jacobi operators (see, e.g., [17, 13] and [9, 2, 7] and references therein).

The results of Sections 6 and 7 were inspired by [3, 4]. We use a different method and obtain more general results. The benefit of our approach is in that:

- we can treat $l > 0$;
- we can insert eigenvalues in the gaps;
- the positivity constraint on the weight can be overcome;

- the proof is very simple and technically undemanding.

Note that the double commutation method of Gesztesy and Teschl [5] gives more or less explicit formula for the Jacobi parameters of the measure with added mass point. Their results work under great generality, of course. However in order to measure the exponential rate of perturbation, as we do here, one would need a good control over the Jost solution.

The only non-elementary tools that we use for the proofs here are Lemmas 2.5 and 2.6 from the author's [10].

Note that these lemmas require all the gaps to be open (see Section 2). Therefore all the results in this paper also inherit this restriction (except Theorems 3.2–3.3 which can be proved using [10, Lemma 4.8]). As soon as Lemmas 2.5 and 2.6 get generalized to the closed gaps situation though, then so will be everything else in this paper.

2. Preliminaries

For the background and additional information on periodic Jacobi matrices, see, e.g. [15]. We follow closely their notation.

For a measure μ with (1.5), the m-function (1.6) is a meromorphic on $\mathbb{C}\setminus\mathfrak{e}$ function. It is Herglotz, meaning $\operatorname{Im} m(z)>0$ whenever $\operatorname{Im} z>0$, and $\operatorname{Im} m(z)<0$ whenever $\operatorname{Im} z<0$. We will discuss its meromorphic continuations through the bands $\mathfrak{e}$ to the second sheet of the Riemann surface $\mathcal{S}_\mathfrak{e}$.

Definition 2.1. *Assume $\mathfrak{e}$ is a finite gap set* (1.5). *Define $\mathcal{S}_\mathfrak{e}$ to be the be the hyperelliptic Riemann surface corresponding to the polynomial $\prod_{j=1}^{l+1}(z-\alpha_j)(z-\beta_j)$.*

We will not give the formal definition, which can be found in many textbooks (see, e.g., [15, Sect 5.12]). Informally $\mathcal{S}_\mathfrak{e}$ can be described as follows.

Let $\mathbb{C}_+=\{z:\operatorname{Im} z>0\}$, $\mathbb{C}_-=\{z:\operatorname{Im} z<0\}$. Denote $\mathcal{S}_+$ and $\mathcal{S}_-$ to be two copies of $\mathbb{C}\cup\{\infty\}$ with a slit along $\mathfrak{e}$ (include $\mathfrak{e}$ as a top edge and exclude it from the lower), and let $\mathcal{S}_\mathfrak{e}$ be $\mathcal{S}_+$ and $\mathcal{S}_-$ glued together along $\mathfrak{e}$ in the following way: passing from $\mathbb{C}_+\cap\mathcal{S}_+$ through $\mathfrak{e}$ takes us to $\mathbb{C}_-\cap\mathcal{S}_-$, and from $\mathbb{C}_-\cap\mathcal{S}_+$ to $\mathbb{C}_+\cap\mathcal{S}_-$. It is clear that topologically $\mathcal{S}_\mathfrak{e}$ is an orientable manifold of genus l.

Let $\pi:\mathcal{S}\to\mathbb{C}\cup\{\infty\}$ be the "projection map" which extends the natural inclusions $\mathcal{S}_+\hookrightarrow\mathbb{C}\cup\{\infty\}$, $\mathcal{S}_-\hookrightarrow\mathbb{C}\cup\{\infty\}$.

The following notation will be used frequently throughout the paper.

Definition 2.2.

- *For $z\in\mathbb{C}\cup\{\infty\}$, denote by z_+ and z_- the two preimages $\pi^{-1}(z)$ in $\mathcal{S}_+$ and $\mathcal{S}_-$ respectively* (*for $z\in\cup_{j=1}^{l+1}\{\alpha_j,\beta_j\}$, z_+ and z_- coincide*).
- *Let $z^\sharp$ be $\left(\overline{\pi(z)}\right)_-$ if $z\in\mathcal{S}_+\setminus\pi^{-1}(\mathfrak{e})$, and $\left(\overline{\pi(z)}\right)_+$ if $z\in\mathcal{S}_-\setminus\pi^{-1}(\mathfrak{e})$. In order to make this continuous, we make the convention $z^\sharp=z$ for $z\in\pi^{-1}(\mathfrak{e})$.*
- *Let $m^\sharp(z)=\overline{m(z^\sharp)}$.*

Note that in fact $m(\bar z)=\overline{m(z)}$ for $z\in\mathcal{S}_+$. In (3.1) we will see that $m(\bar z)=\overline{m(z)}$ for $z\in\mathcal{S}_-$ as well, so one can think of $m^\sharp$ as $m((\pi(z))_-)$ if $z\in\mathcal{S}_+$ and $m((\pi(z))_+)$ if $z\in\mathcal{S}_-$.

As was mentioned in the Introduction, the essential spectrum of any p-periodic Jacobi matrix is a finite gap set (1.5), where $l\le p-1$. In fact, there exists a polynomial Δ (called the discriminant) of degree p such that

$$\mathfrak{e} = \bigcup_{j=1}^{l+1} [\alpha_j, \beta_j] = \Delta^{-1}([-2,2]).$$

In particular $\Delta(z) = z$ when $\mathfrak{e} = [-2,2]$.

If $l = p-1$, then we say that $\mathcal{J}$ has all gaps open. It is known that a finite gap set $\mathfrak{e}$ is an essential spectrum of some periodic Jacobi matrix if and only if the harmonic measure of each band is rational. It is an essential spectrum of some periodic Jacobi matrix with all gaps open if and only if the harmonic measure of each band is equal.

It turns out that if there exists at least one periodic Jacobi matrix $\mathcal{J}$ with $\sigma_{ess}(\mathcal{J}) = \mathfrak{e}$, then there exists a whole set of periodic Jacobi matrices satisfying the same property. In fact, this set is homeomorphic to $(S^1)^g$, a g-dimensional torus. See [15, Chapt 5] for more details. This motivates the following definition.

Definition 2.3. *The isospectral torus* $\mathcal{T}_{\mathfrak{e}}$ *of* $\mathfrak{e}$ *is the set of periodic Jacobi matrices* $\mathcal{J}$ *with* $\sigma_{ess}(\mathcal{J}) = \mathfrak{e}$*.*

From now on assume that all gaps of $\mathfrak{e}$ are open (this is a generic situation, see [15, Chapt 5]).

Let us define some special subsets $\mathcal{S}_R$ of $\mathcal{S}_{\mathfrak{e}}$ that will be used commonly in this paper.

Definition 2.4. *Let* $x(z) = z + z^{-1}$*. For each* $R > 1$*, let*

$$\mathcal{S}_R = \mathcal{S}_+ \cup \pi^{-1}(E_R), \tag{2.1}$$

where E_R *is the union of interiors of the bounded components of the set* $\Delta^{-1}(x(R\,\partial\mathbb{D}))$*.*

Note that $x(R\,\partial\mathbb{D})$ are concentric ellipses. Below is an example ($p = 3$) how E_R evolves as R grows.



Now we are ready to state the main tools of this paper.

Lemma 2.5 ([10]). *Assume* $\operatorname{ess\,supp} \mu = \mathfrak{e}$*, and let* $m(z) = \int \frac{d\mu(x)}{x-z}$*.*

The following are equivalent:

(i) The Jacobi matrix $(a_n, b_n)_{n=1}^{\infty}$ *associated with* μ *satisfies*

$$\limsup_{n\to\infty} \left(|a_n - a_n^{(0)}| + |b_n - b_n^{(0)}| \right)^{1/2n} \leq R^{-1}, \tag{2.2}$$

where $(a_n^{(0)}, b_n^{(0)})_{n=1}^{\infty}$ *is a periodic Jacobi matrix from* $\mathcal{T}_{\mathfrak{e}}$*.*

(ii) (a) *m has a meromorphic continuation to* $\mathcal{S}_R$*;*

(b) *m has no poles on $\pi^{-1}(\mathfrak{e})$, except at $\pi^{-1}(\cup_{j=1}^{p}\{\alpha_j,\beta_j\})$, where they are at most simple;*
(c) *$m(z)-m^\sharp(z)$ has no zeros in $\pi^{-1}(E_R)$, except at $\pi^{-1}(\cup_{j=1}^{p}\{\alpha_j,\beta_j\})$, where they are at most simple;*
(d) *If m has a pole at z for $z\in\pi^{-1}(E_R\setminus\mathfrak{e})$ then $z^\sharp$ is not a pole of m.*

Lemma 2.6 ([10])**.** *Assume* $\operatorname{ess\,supp}\mu=\mathfrak{e}$*, and let* $m(z)=\int\frac{d\mu(x)}{x-z}$.
The following are equivalent:

(i) The Jacobi matrix $(a_n,b_n)_{n=1}^\infty$ associated with μ is eventually periodic, i.e., satisfies

$$(a_n,b_n)_{n=s}^\infty\in\mathcal{T}_{\mathfrak{e}}\quad\text{for some } s. \tag{2.3}$$

(ii) (a) *m has a meromorphic continuation to $\mathcal{S}$;*
(b) *m has no poles on $\pi^{-1}(\mathfrak{e})$, except at $\pi^{-1}(\cup_{j=1}^{p}\{\alpha_j,\beta_j\})$, where they are at most simple;*
(c) *$m(z)-m^\sharp(z)$ has no zeros in $\mathcal{S}\setminus\{\pm\infty\}$, except at $\pi^{-1}(\cup_{j=1}^{p}\{\alpha_j,\beta_j\})$, where they are at most simple;*
(d) *If m has a pole at z for $z\in\pi^{-1}(\mathbb{C}\setminus\mathfrak{e})$ then $z^\sharp$ is not a pole of m.*

Because we will be discussing meromorphic continuations, the next lemma will prove to be useful.

Lemma 2.7 ([6])**.** *Let m be as in* (1.6)*. Then m can be analytically continued from $\mathcal{S}_+\cap\pi^{-1}(\mathbb{C}_+)$ through an interval $I\subset\mathbb{R}$ if and only if the associated measure μ is purely absolutely continuous on I, and the density $f(x)=\frac{d\mu}{dx}$ is real-analytic on I. In this case, the analytic continuation of m into some domain $\mathcal{D}_-$ of $\mathcal{S}_-\cap\pi^{-1}(\mathbb{C}_-)$ is given by*

$$m(z_-)=\overline{m(\bar z_+)}+2\pi i f(z),\quad z\in\pi(\mathcal{D}_-),$$

where $f(z)$ is the complex-analytic continuation of f to $\mathcal{D}_-$.

Definition 2.8. *Suppose m satisfies either Lemma* (2.5) *or* (2.6)*, and let $z\in\mathcal{S}_{\mathfrak{e}}$ be a pole of m. Then*

- *We call $\pi(z)$ a singularity of $\mathcal{J}$;*
- *If $z\in\mathcal{S}_+\setminus\cup_{j=1}^{l+1}\{\alpha_j,\beta_j\}$, then we say that $\pi(z_j)$ is an eigenvalue of $\mathcal{J}$;*
- *If $z\in\mathcal{S}_-$, then we say that $\pi(z_j)$ is a resonance of $\mathcal{J}$;*
- *If $z\in\mathcal{S}_-$ and $\operatorname{Im}\pi(z)=0$, then we say that $\pi(z_j)$ is an antibound state of $\mathcal{J}$.*

Note that a band edge $z_0\in\{\alpha_j,\beta_j\}$ may be a pole of m (which means $m(z_0)\sim O(\frac{1}{(z-z_0)^{1/2}})$), in which case it is a resonance, as well as an antibound state.

As a notational remark, whenever we have any function g of complex variable, and $z\in\mathcal{S}$, then we will occasionally write $g(z)$, instead of $g(\pi(z))$.

3. Direct Problem

3.1. Structure of singularities of exponentially small perturbations of periodic Jacobi matrices.

Suppose $\mathcal{J}$ satisfies (1.4) for some $R>1$. Note that for $x\in\mathfrak{e}$,

$$\operatorname{Im} m((x+i0)_+)=\pi f(x)=-\operatorname{Im} m((x-i0)_+).$$

Thus using Lemma 2.7 to continue m from $\mathcal{S}_+\cap\pi^{-1}(\mathbb{C}_+)$ to $\mathcal{S}_-\cap\pi^{-1}(\mathbb{C}_-)$, and then from $\mathcal{S}_+\cap\pi^{-1}(\mathbb{C}_-)$ to $\mathcal{S}_-\cap\pi^{-1}(\mathbb{C}_+)$, we obtain that for $z\in\mathcal{S}_-\cap\pi^{-1}(\mathbb{C}_+)\cap\mathcal{S}_R$,

$$m(z_-) = m(z_+) + 2\pi i f(z),$$
$$m(\bar z_-) = m(\bar z_+) - 2\pi i f(z),$$

which, using $\overline{m(z_+)} = m(\bar z_+)$, implies

$$\overline{m(z_-)} = m(\bar z_-), \quad z\in\mathcal{S}_-\cap\pi^{-1}(\mathbb{C}_+)\cap\mathcal{S}_R. \tag{3.1}$$

Now we establish properties of singularities (see Definition 2.8) of $\mathcal{J}$. The main condition is (iv), which, roughly speaking, says that every second singularity of m must be necessarily a resonance when counted either from the left edge of any band or from the right.

Theorem 3.1. *Suppose $\mathcal{J}$ satisfies* (1.4) *for some $1 < R \le \infty$. Denote by $\{z_j\}_{j=1}^N$ ($N \le \infty$) the singularities of m (counting multiplicities) that are located in the simply-connected component of $E_R\cap\mathbb{R}$ which contains $[\alpha_j,\beta_j]$. Denote by $x_1, x_2, \dots$ and $y_1, y_2, \dots$ to be the real singularities of m in that region:*

$$\dots \le y_2 \le y_1 \le \alpha_j < \beta_j \le x_1 \le x_2 \le \dots$$

Then

- (i) *$z_j \in \mathbb{C}\setminus\cup_{j=1}^{l+1}(\alpha_j,\beta_j)$;*
- (ii) *Non-real z_j's come is complex conjugate pairs* (*counting multiplicities*)*, and are necessarily resonances;*
- (iii) *If $x_1 = \beta_j$ then x_1 is a resonance of order 1; if $y_1 = \alpha_j$ then y_1 is a resonance of order 1;*
- (iv) *$x_2, x_4, x_6, \dots$ and $y_2, y_4, y_6 \dots$ are antibound states of m.*

Remark. In particular, (iv) implies that:

- Between any two eigenvalues (which are located in the same simply-connected component of E_R and are not separated by a band) there is an odd number of resonances (counting multiplicities);
- If a singularity is of order higher than 1, then it's necessarily a resonance.

Proof. (i) and (iii) follow from Lemma 2.5(ii)(b). (ii) follows from the fact that m is Herglotz and (3.1). Let us prove (iv) now.

Since m has at most order 1 pole at band edges, let the Taylor's series of m at β_j be

$$m(z) = c_{-1}(z-\beta_j)^{-1/2} + c_0 + c_1(z-\beta_j)^{1/2} + c_2(z-\beta_j) + \dots$$

Since m is Herglotz, $\operatorname{Im} m(x+i0) \ge 0$ on $[\alpha_j,\beta_j]$. This implies that either $c_{-1} < 0$, or $c_{-1} = 0$ and $c_1 > 0$ (note that it is not possible to have $c_{-1} = c_1 = 0$ because zeros of $m - m^\sharp$ at the band edges are at most simple by Lemma 2.5(ii)(c)). Then

$$m^\sharp(z) = -c_{-1}(z-\beta_j)^{-1/2} + c_0 - c_1(z-\beta_j)^{1/2} + c_2(z-\beta_j) + \dots,$$

and thus

$$m(z) - m^\sharp(z) = 2c_{-1}(z-\beta_j)^{-1/2} + 2c_1(z-\beta_j)^{1/2} + O((z-\beta_j)^{3/2}).$$

Now, if $c_{-1} < 0$, then $m - m^\sharp$ is negative to the right of β_j. Since in this case β_j is a first order pole, $x_1 = \beta_j$. Note that $m - m^\sharp$ cannot be equal to zero

(Lemma 2.5(ii)(c)) on (β_j, α_{j+1}), and thus

$$\lim_{z\to(x_2-0)_+} m(z) - m^\sharp(z) = -\infty \tag{3.2}$$

This implies that x_2 must be a resonance: indeed, if $(x_2)_+$ were an eigenvalue, then $\lim_{z\to(x_2-0)_+} m(z) = +\infty$ (because $\lim_{x\to x_2-0} \frac{1}{x_2-x} = +\infty$) and $\lim_{z\to(x_2-0)_+} m^\sharp(z)$ is finite (Lemma 2.5(ii)(d)), which would contradict to (3.2).

Now, if $c_{-1} = 0$ and $c_1 > 0$, then $m - m^\sharp$ is positive to the right of β_j. Since $m - m^\sharp$ cannot be equal to zero, we obtain

$$\lim_{z\to(x_1-0)_+} m(z) - m^\sharp(z) = +\infty.$$

If $x_1 = x_2$ then it is a resonance, since m is Herglotz on $\mathcal{S}_+$ and therefore cannot have second order poles there. If $x_1 \neq x_2$, then

$$\lim_{z\to(x_1+0)_+} m(z) - m^\sharp(z) = -\infty,$$
$$\lim_{z\to(x_2-0)_+} m(z) - m^\sharp(z) = -\infty,$$

which again implies that x_2 is a resonance by the arguments above. That x_k is a resonance for every even k follows by the same arguments.

Similarly for y_j's: if the Taylor's series of m at α_j is

$$m(z) = c_{-1}(z-\alpha_j)^{-1/2} + c_0 + c_1(z-\alpha_j)^{1/2} + c_2(z-\alpha_j) + \dots,$$

then in order to have $\operatorname{Im} m(x+i0) \geq 0$ on $[\alpha_j, \beta_j]$, we must have that c_{-1} is purely imaginary with $\operatorname{Im} c_{-1} > 0$, or $c_{-1} = 0$ and c_1 is purely imaginary with $\operatorname{Im} c_1 > 0$. Then

$$m(z) - m^\sharp(z) = 2c_{-1}(z-\alpha_j)^{-1/2} + 2c_1(z-\alpha_j)^{1/2} + O((z-\alpha_j)^{3/2}).$$

In the first case scenario, $\operatorname{Im} c_{-1} > 0$, $y_1 = \alpha_j$, $m - m^\sharp$ positive to the left of α_j, so

$$\lim_{z\to(y_2+0)_+} m(z) - m^\sharp(z) = +\infty, \tag{3.3}$$

which implies that y_2 is a resonance. In the second case, $c_{-1} = 0$, $\operatorname{Im} c_1 > 0$, $m - m^\sharp$ is negative to the left of α_j, so we obtain

$$\lim_{z\to(y_1+0)_+} m(z) - m^\sharp(z) = -\infty,$$
$$\lim_{z\to(y_1-0)_+} m(z) - m^\sharp(z) = +\infty,$$
$$\lim_{z\to(y_2+0)_+} m(z) - m^\sharp(z) = +\infty,$$

which again implies that y_2 is a resonance. □

3.2. **Eventually periodic situation.**

Theorem 3.2. *Let $\mathcal{J}$ be eventually periodic, that is* (1.3)*, and let m be its m-function* (1.6)*. Then*

$$m(z) = \frac{p(z) \pm \sqrt{r(z)}}{a(z)},$$
$$m(z) \sim -\frac{1}{z}, \quad z \to \infty_+,$$

where

$$r(z) = \prod_{j=1}^{l+1} (z-\alpha_j)(z-\beta_j), \tag{3.4}$$

$$a(z) = A\prod_{j=1}^{N} (z-z_j), \tag{3.5}$$

$$\deg a = N \geq l, \tag{3.6}$$

$$\deg p \leq \max\{N-1, l+1\}. \tag{3.7}$$

The singularities $\{z_j\}_{j=1}^N$ ($N<\infty$) *of* m *satisfy the following properties:*

(i) $z_j \in \mathbb{C} \setminus \cup_{j=1}^{l+1}(\alpha_j, \beta_j)$*;*

(ii) *Non-real* z_j*'s come is complex conjugate pairs (counting multiplicities), and are necessarily resonances;*

(iii) *If* z_j *is a band edge then it is a resonance of multiplicity* 1*;*

(iv) *When real singularities are counted (with multiplicities) from any band edge (as described in Thm 3.1), then every even-numbered one is necessarily a resonance.*

Remarks. 1. In particular, (iv) implies that:

- Between any two eigenvalues (which are not separated by a band) there is an odd number of resonances (counting multiplicities);
- Each gap $[\beta_j, \alpha_{j+1}]$ contains odd number of real singularities (counting multiplicities);
- If a singularity is of order higher than 1, then it's necessarily a resonance.

2. In the proof of this Theorem we use Lemma 2.6(i)⇒(ii), which demands all gaps to be open. However, the direction Lemma 2.6(i)⇒(ii) can be established easily in the closed gap situation via induction and [10, Lemma 4.8]).

3. If $\mathcal{J}$ is actually periodic, then (i) should say $z_j \in \mathbb{C}\cup\{\infty\} \setminus \cup_{j=1}^{l+1}(\alpha_j, \beta_j)$, since ∞_- is actually a first pole of m. However if $\mathcal{J}$ is not periodic, then $\infty_\pm$ is never a pole, as we show in Theorem 3.3.

Proof. By [15, Prop. 5.12.1], any m meromorphic on $\mathcal{S}$ must be of the form

$$m(z) = \frac{p(z) \pm q(z)\sqrt{r(z)}}{a(z)},$$

where p, q, a are polynomials with no common zeros, and r as in (3.4). However by Lemma 2.6(ii)(c), q cannot have zeros anywhere except possibly at the band edges. Suppose $q(\alpha_j) = 0$. Lemma 2.6(ii)(c) states that $m(z) - m^\sharp(z) = \frac{2q(z)\sqrt{r(z)}}{a(z)}$ has at most first order zero at α_j. Therefore α_j must also be a zero of $a(z)$. But then $p(\alpha_j) \neq 0$, which implies that $m = O\left(\frac{1}{(z-\alpha_j)^k}\right)$ with $k \geq 1$, which is a pole of order at least 2 at α_j. This contradiction shows that q is a constant and therefore may be assumed to be 1.

The degree of m as a meromorphic function on $\mathcal{S}_\mathfrak{e}$ (i.e. topological map from $\mathcal{S}_\mathfrak{e}$ into the Riemann sphere) is at least $l+1$: see [15, Thm 5.12.5(b)]. Thus it has at least $l+1$ poles. Therefore either the degree N of the polynomial a is at least l, or otherwise ∞_- would be a pole of order 2 (∞_+ is a zero of m so cannot be a pole). However ∞_- is never pole of order 2: it is either a pole of order 1 if the Jacobi matrix is periodic (see [15, Thm 5.13.1(iv)]), and otherwise it is not a pole (as we show in Thm 3.3 below).

This proves that $\deg a \geq l$. Note that at ∞_+, $\sqrt{r(z)} \sim z^{l+1}$, so in order to have $m(z) \sim -\frac{1}{z}$ at ∞_+ we must have: if $\deg a = l$ then $\deg p = l+1$; if $\deg a = l+1$ or $l+2$ then $\deg p \leq l+1$; if $\deg a > l+2$ then $\deg p = \deg a - 1$.

(i) follows from Lemma 2.6(ii)(b).

(ii) follows from the fact that m is Herglotz on $\mathcal{S}_+$ and (3.1).

(iii) and (iv) follow from Theorem 3.1. ☐

Let us introduce the notation

$$\mathcal{J}^{(s)} = (a_n, b_n)_{n=s+1}^{\infty}, \tag{3.8}$$

that is, $\mathcal{J}^{(s)}$ is the Jacobi matrix obtained from $\mathcal{J}$ by removing the first s rows and columns. It is well-known that the m-functions $m(z)$, $m^{(1)}(z)$ of $\mathcal{J}$ and $\mathcal{J}^{(1)}$ obey

$$m(z) = \frac{1}{b_1 - z - a_1^2 m^{(1)}(z)}. \tag{3.9}$$

Suppose $\mathcal{J}$ is an eventually periodic Jacobi matrix with (1.3) for some $s \geq 0$. This says that $\mathcal{J}^{(s)} \in \mathcal{T}_{\mathfrak{e}}$. Suppose $\mathcal{J}^{(s-1)} \notin \mathcal{T}_{\mathfrak{e}}$. We will say that coefficient a_s is aperiodic if $a_s \neq a_{s+p}$, and otherwise we will call it periodic. Similarly for b_s. Since $\mathcal{J}^{(s-1)} \notin \mathcal{T}_{\mathfrak{e}}$, then either a_s, or b_s, or both of them are aperiodic.

Theorem 3.3. *Suppose $\mathcal{J}$ is an eventually periodic Jacobi matrix with $\mathcal{J}^{(s)} \in \mathcal{T}_{\mathfrak{e}}$ but $\mathcal{J}^{(s-1)} \notin \mathcal{T}_{\mathfrak{e}}$ ($s \geq 0$). Then:*

- (i) *If $s = 0$ then $\deg m = l+1$ and the $l+1$ singularities of $\mathcal{J}$ are: one at ∞_- and one per each gap.*
- (ii) *If $s > 0$, b_s is aperiodic, and a_s is periodic, then $\deg m = l + 2s - 1$, and ∞_- is not a pole of m.*
- (iii) *If $s > 0$ and a_s is aperiodic, then $\deg m = l + 2s$, and ∞_- is not a pole of m.*

Proof. For the periodic case $s = 0$ the result is proved in [15, Thm 5.13.1].

Now suppose $s = 1$. Then

$$m(z) = \frac{1}{b_1 - z - a_1^2 m^{(1)}(z)}, \tag{3.10}$$

where $m^{(1)}$ has one singularity per gap and one at ∞_-. Recall that $m^{(1)}(z) \sim -\frac{1}{z}$ at ∞_+ and $m^{(1)}(z) \sim k_1 z + k_0 + O(\frac{1}{z})$, $k_1 \neq 0$, at ∞_-. Thus $m^{(1)}(z) = 0$ exactly once per each gap, at ∞_+, and possibly at ∞_-. Note that $m^{(1)}(z) = 0$ at ∞_- if and only if $-1 - a_1^2 k_1 \neq 0$. But we know that if a_1 is periodic then $m^{(1)}(z)$ has a pole at ∞_-. Therefore $a_1 = \sqrt{-1/k_1}$ is precisely the condition for a_1 to be periodic.

Thus, we have shown that if $s = 1$ and a_1 is aperiodic then m has exactly $l+2$ zeros: once per each gap, one at ∞_+, and one at ∞_-. Thus $\deg m = l+2$.

Suppose now that a_1 is periodic, $a_1 = \sqrt{-1/k_1}$, but b_1 is aperiodic. Then ∞_- is not a zero of m, so $\deg m = l+1$. Moreover, note that ∞_- is a pole of m if and only if $b_1 - a_1^2 k_0 = 0$. This means that $b_1 + k_0/k_1 = 0$ is exactly the condition for b_1 to be periodic. Thus, if $s = 1$, a_1 is periodic, and b_1 is aperiodic, then $\deg m = l+1$, and ∞_- is not a pole of m.

Now $s \geq 2$ follows easily by induction. Indeed,

$$m(z) = \frac{1}{b_1 - z - a_1^2 m^{(1)}(z)}, \tag{3.11}$$

and ∞_- is not a pole of $m^{(1)}$. Therefore m has zeros at ∞_+, ∞_-, and at every pole of $m^{(1)}$. Therefore $\deg m = \deg m^{(1)} + 2$. $\square$

4. Inverse problem

Theorem 4.1. *Let* $\{(z_j, \epsilon_j)\}_{j=1}^N$ $(l+1 \leq N < \infty)$ *be given* (*not necessarily distinct*), *where* $z_j \in \mathbb{C}$, $\epsilon_j \in \{1, -1\}$. *If* $\epsilon_j = 1$ *then we think of* z_j *as an eigenvalue, and if* $\epsilon_j = -1$ *then as a resonance.*

If this set of point satisfies (i)–(iv) of Theorem 3.2, then there exists a unique eventually periodic Jacobi matrix $\mathcal{J}$ *with such a system of singularities.*

Moreover, if $s = \lceil \frac{N-l}{2} \rceil$, *then* $\mathcal{J}^{(s)}$ *is periodic, and* $\mathcal{J}^{(s-1)}$ *is not. If* $N-l$ *is odd then then* a_s *is periodic and* b_s *is aperiodic; if* $N-l$ *is even then* a_s *is aperiodic.*

Remarks. 1. If $N = l+1$, but not all of them are finite (namely, $(z_{N+1}, \epsilon_{N+1}) = (\infty_-, -1)$, a resonance at ∞_-), then there exists a unique periodic Jacobi matrix with such a system of singularities. This is well known, see [15, Thm 5.13.5] and references therein.

2. In particular, we implicitly assume that our system $\{(z_j, \epsilon_j)\}_{j=1}^N$ satisfies the conditions in Remark 1 after Theorem 3.2 (which are just corollaries of (iv)).

3. As becomes clear from the proof, the spectral measure of any eventually periodic Jacobi matrix can be explicitly written in terms of the system of singularities, see (4.4)–(4.6).

Proof. Let

$$m(z) = \frac{p(z) + \sqrt{r(z)}}{a(z)}, \quad z \in \mathcal{S}_+, \tag{4.1}$$

where $r(z)$ is (3.4), $a(z) = A \prod_{j=1}^N (z - z_j)$, and A, p are to be determined. Note that by Lemma 2.6(ii)(d), we want to avoid simultaneous poles of m at $(z_j)_+$ and $(z_j)_-$. Therefore we must require

$$p(z_j) = \epsilon_j \sqrt{r(z_j)}, \quad j = 1, \ldots, N. \tag{4.2}$$

However, if z_j is a singularity of order $s \geq 2$, i.e., $z_j = z_{j+1} = \ldots = z_{j+s-1}$, then additionally we require

$$p^{(m)}(z_j) = -\frac{d^m}{dz^m} \sqrt{r(z)}\Big|_{z=z_j}, \quad m = 0, 1, \ldots, s-1. \tag{4.3}$$

Here $\epsilon_j = -1$ since it is implied by the property (iv) of Theorem 3.2.

Note that (4.2)+(4.3) form a system of N equations (if we ignore the identical equations in (4.2)) with respect to the unknown coefficients of the polynomial

$$p(z) = \sum_{k=0}^M c_k z^k.$$

It is easy to see that if the number of unknowns equals to the number of equations that this system has a unique solution. We show this in the next lemma.

Lemma 4.2. *If* $M = N - 1$ *then the system* (4.2)+(4.3) *has a unique solution.*

Proof. If all z_j's have multiplicity 1, then the coefficient matrix of our linear system is a Vandermonde matrix, and therefore has a non-vanishing determinant.

If some of the z_j's have multiplicity ≥ 2, then one needs to slightly modify the standard proof of Vandermonde determinant. Indeed, performing the elementary row operations the determinant of the coefficient matrix can be reduced to

$$\pm \prod_{j<k; z_j \neq z_k} (z_j - z_k)^{s_j s_k},$$

where s_j is the multiplicity of z_j. $\square$

Now, let us consider the following cases.

If $N = l+1$ then we take $M = l+1$, $c_{l+1} = -1$ (this ensures that $m(z) \to 0$ at ∞_+), and the rest of the coefficients $c_0, \dots, c_l$ to be the unique solution of (4.2)+(4.3) (Lemma 4.2 applies since we can take the c_{l+1}-terms to the right-hand side, and therefore the number of unknowns equals to the number of equations).

If $N \geq l+2$ then we take $M = N-1$, and choose $c_0, \dots, c_M$ to be the unique solution of (4.2)+(4.3).

Now for any $z \in \mathcal{S}_+ \cap \pi^{-1}(\mathbb{C}_+)$, let us choose Γ_+ to be any simple clockwise contour in $\mathcal{S}_+$ whose interior contains $\mathfrak{e}$, all the singularities $\{z_j\}_{j=1}^N$, but does not contain z. Note that by our construction above, $m(z) \to 0$ as $z \to \infty_+$. Thus by the Cauchy formula,

$$\frac{1}{2\pi i} \int_{\Gamma_+} \frac{m(w)}{w - z} dw = m(z), \quad z \in \mathcal{S}_+.$$

Now by shrinking the contour towards the real axis, and using (4.1) and (4.2), we obtain that

$$m(z) = \int_{\mathfrak{e}} \frac{g(x)}{x - z} + \sum_{\{j : \epsilon_j = 1\}} \frac{w_j}{z_j - z}, \quad z \in \mathcal{S}_+, \tag{4.4}$$

where

$$g(x) = \frac{1}{\pi} \frac{\lim_{\varepsilon \to 0+} \operatorname{Im} \sqrt{r(x + i\varepsilon)}}{A \prod_{k=1}^N (x - z_k)} \tag{4.5}$$

$$w_j = -\frac{2\sqrt{r(z_j)}}{A \prod_{k \neq j} (z_j - z_k)} \tag{4.6}$$

Note that $\sqrt{r(z)}$ here switches sign from one gap to another, and $\lim_{\varepsilon \to 0+} \operatorname{Im} \sqrt{r(x + i\varepsilon)}$ switches sign from one band to another.

We still have not defined A. First choose the sign of A to be such that (4.5) is positive for $x \in (\alpha_{l+1}, \beta_{l+1})$. Note that the sign of $\lim_{\varepsilon \to 0+} \operatorname{Im} \sqrt{r(x + i\varepsilon)}$ on (α_l, β_l) is opposite to its sign on $(\alpha_{l+1}, \beta_{l+1})$. However a has odd number of roots on $[\beta_l, \alpha_{l+1}]$, which means that $g(x)$ is also positive on (α_l, β_l). Proceeding further in the same manner, one shows that $g(x) \geq 0$ on $\mathfrak{e}$.

Now let us show that all the weights w_j are positive. Suppose z_{j_0} is the smallest singularity on $[\alpha_{l+1}, \infty)$. Suppose that $\epsilon_{j_0} = 1$. Then the sign of w_{j_0}, see (4.6), coincides with the sign of g on $(\alpha_{l+1}, \beta_{l+1})$, see (4.5), i.e., is positive. Similarly, every odd-numbered w_j on (α_{l+1}, ∞) is positive. Tracking signs in other gaps and on $(-\infty, \alpha_1]$ in the similar fashion proves that all w_j (with $\epsilon_j = 1$) are positive.

Therefore (4.4) implies that m is a Herglotz function:

$$m(z) = \frac{d\mu(x)}{x - z}$$

for some positive finite measure μ. Now we can choose the absolute value of A so that μ has total mass 1.

This shows that m is an m-function of some Jacobi matrix $\mathcal{J}$. By the construction, it is clear that m satisfies all the conditions of Lemma 2.6(ii). Thus Lemma 2.6 proves that $\mathcal{J}$ is eventually periodic.

Uniqueness is clear: each step of the construction was uniquely determined by the requirements of Theorem 3.2.

The "moreover" part of Theorem 4.1 immediately follows from Theorem 3.3. $\square$

Corollary 4.3. *Let S be a finite gap set $\mathfrak{e}$ with finitely many points outside of it. Then there exists an eventually periodic Jacobi matrix $\mathcal{J}$ with*

$$\sigma(\mathcal{J}) = S.$$

Remarks. 1. Given the current limitations of Lemma 2.6, $\mathfrak{e}$ must consist of bands of equal equilibrium measure.

2. Of course such $\mathcal{J}$ is far from being unique because we can choose resonances in infinitely many ways.

5. $l = 0$ AND PERTURBATION DETERMINANTS

Using the above ideas (for $l = 0$), one can now restate the results of Damanik–Simon, [1], in a more explicit way. Recall that in the perturbations of the free case, under sufficient regularity on the Jacobi parameters of $\mathcal{J}$ (e.g. if the Jacobi parameters obey $\sum_{j=1}^{\infty} |a_j|^2 + |1 - b_j|^2 < \infty$), one can define the perturbation determinant of $\mathcal{J}$ via

$$L(z) = \det\Big(1 + (\mathcal{J} - \mathcal{J}_0)\left[\mathcal{J}_0 - (z + z^{-1})\right]^{-1}\Big)$$

(see [1, 8]). Another way to think about it is as $L(z) = \frac{u(z)}{u(0)}$, where $u(z)$ is the Jost function of $\mathcal{J}$. Note that $u(0) > 0$ (see [1, Eq. (A16)]. The main results of [1] are the two theorems ([1, Thm 1.9 and Thm 1.11]) formulated below. To state them we need the notion of canonic weights from [1]. This notion is not the most straightforward and is indirect in terms of L. Even in terms of u, there is an implicit condition of positivity of the right-hand side of (5.1). The main purpose of the current section is to show that canonicity can be expressed more directly.

Definition 5.1. *Suppose a function u is real analytic in $\{z : |z| < R\}$ for some $R > 1$, has only real zeros $\{z_j\}_{j=1}^n$ in $\mathbb{D}$, and let a set of positive numbers $\{w_j\}_{j=1}^n$ be given.If $|z_j| > R^{-1}$, then w_j is said to be canonical if and only if*

$$w_j = \frac{(z_j - z_j^{-1})^2}{z_j}\left[u'(z_j)\overline{u(1/\bar{z}_j)}\right]^{-1} \tag{5.1}$$

Theorem 5.2 (Damanik–Simon)**.** *If $a_n = 1$ and $b_n = 0$ for large n, then $L(z)$ is a real polynomial and all the weights are canonical. Conversely, if $L(z)$ satisfies*

(i) *$L(z)$ is a real polynomial;*
(ii) *$L(z)$ is nonvanishing on $\overline{\mathbb{D}} \setminus \mathbb{R}$;*
(iii) *if $+1$ and/or -1 are zeros, they are simple;*

(iv) $L(0) = 1$,

then there is at most one set of Jacobi parameters with $a_n = 1$ *and* $b_n = 0$ *for* n *large that has* L *as perturbation determinant. Moreover, the weights associated to this set are the canonical ones. If these canonical weights lead to* $w_j > 0$*, then there is a set of Jacobi parameters with* $a_n = 1$ *and* $b_n = 0$ *for large* n.

Theorem 5.3 (Damanik–Simon)**.** *Let* $R > 1$*. If*

$$\limsup_{n\to\infty} \left(|1 - a_n| + |b_n|\right)^{1/2n} \le R^{-1} \tag{5.2}$$

holds, then $L(z)$ *has a real analytic continuation to* $\{z : |z| < R\}$ *and the weights of all* z_j *with* $|z_j| > R^{-1}$ *are canonical. Conversely, if* $L(z)$ *is real analytic in* $\{z : |z| < R\}$ *and obeys (ii)–(iv) of Theorem 5.2, then* (5.2) *holds if and only if all weights for* z_j *with* $|z_j| > R^{-1}$ *are canonical.*

Below we reformulate these theorems in terms of properties of L alone, and the notion of canonicity can be avoided.

Theorem 5.4. $L(z)$ *is a perturbation determinant of an eventually free Jacobi matrix if and only if:*

- (i) $L(z)$ *is a real polynomial;*
- (ii) $L(z)$ *is nonvanishing on* $\overline{\mathbb{D}} \setminus \mathbb{R}$*;*
- (iii) *if* $+1$ *and/or* -1 *are zeros, they are simple;*
- (iv) $L(0) = 1$*;*
- (v) *Let* $0 < x_n < \ldots < x_2 < x_1 < 1$ *be the positive zeros of* L *in* $\mathbb{D}$ *counting multiplicities. Then*
  - (a) *There is an even number of zeros (counting multiplicities) of* L *on* $[1, x_1^{-1})$*;*
  - (b) *There is an odd number of zeros (counting multiplicities) of* L *on* (x_j^{-1}, x_{j+1}^{-1}) $(j \ge 1)$*;*
  - (c) x_j^{-1} *is not a zero of* L*;*
- (vi) *Let* $-1 < y_1 < y_2 < \ldots < y_n < 0$ *be the negative zeros of* L *in* $\mathbb{D}$ *counting multiplicities. Then*
  - (a) *There is an even number of zeros (counting multiplicities) of* L *on* $(y_1^{-1}, -1]$*;*
  - (b) *There is an odd number of zeros (counting multiplicities) of* L *on* (y_{j+1}^{-1}, y_j^{-1}) $(j \ge 1)$*;*
  - (c) y_j^{-1} *is not a zero of* L*.*

Theorem 5.5. *Let* $R > 1$*. Then* $L(z)$ *is a perturbation determinant of a Jacobi matrix satisfying*

$$\limsup_{n\to\infty} \left(|1 - a_n| + |b_n|\right)^{1/2n} \le R^{-1} \tag{5.3}$$

if and only if

- (i′) $L(z)$ *has a real analytic continuation to* $\{z : |z| < R\}$*;*
- (ii) $L(z)$ *is nonvanishing on* $\overline{\mathbb{D}} \setminus \mathbb{R}$*;*
- (iii) *if* $+1$ *and/or* -1 *are zeros, they are simple;*
- (iv) $L(0) = 1$*;*
- (v) *Let* $0 < x_n < \ldots < x_2 < x_1 < 1$ *be the positive zeros of* L *in* $\mathbb{D}$ *counting multiplicities, and suppose* $x_{k+1} \le R^{-1} < x_k$*. Then*

   (a) *There is an even number of zeros (counting multiplicities) of* L *on* $[1, x_1^{-1})$*;*
   (b) *There is an odd number of zeros (counting multiplicities) of* L *on* (x_j^{-1}, x_{j+1}^{-1}) $(1 \le j \le k-1)$*;*
   (c) x_j^{-1} *is not a zero of* L $(1 \le j \le k)$*;*

(vi) *Let* $-1 < y_1 < y_2 < \ldots < y_n < 0$ *be the negative zeros of* L *in* $\mathbb{D}$ *counting multiplicities, and suppose* $y_k < -R^{-1} \le y_{k+1}$*. Then*
   (a) *There is an even number of zeros (counting multiplicities) of* L *on* $(y_1^{-1}, -1]$*;*
   (b) *There is an odd number of zeros (counting multiplicities) of* L *on* (y_{j+1}^{-1}, y_j^{-1}) $(1 \le j \le k-1)$*;*
   (c) y_j^{-1} *is not a zero of* L $(1 \le j \le k)$*.*

Proofs of Theorems 5.4 and 5.5. Note that zeros of u coincide with the poles of M, where $M(z) = -m(z+z^{-1})$ for $z \in \{z : |z| < 1\}$. Therefore Theorem 5.4 can be easily deduced by applying Theorems 3.2 and 4.1. But the latter theorem is currently proved only if all the intervals of $\mathfrak{e}$ have equal equilibrium measures. So instead let us prove directly that the existence of canonical weights is equivalent to (v) and (vi).

Note that if all w_j are canonical, then (5.1) implies that $u'(x_j)\overline{u(1/\bar{x}_j)} > 0$ $(j \le k)$.

Suppose first that $u(1) > 0$ or $u(1) = 0$, $u'(1) < 0$. This implies $u'(x_1) > 0$, which means $u(1/x_1) > 0$, i.e., (v)(a) holds. Continuing this further, we get $u'(x_2) < 0$, which then implies $u(1/x_2) < 0$, which means there is an odd number of zeros of u on (x_1^{-1}, x_2^{-1}). This proves (v)(b). (v)(c) is trivial since w_j is finite.

Similarly, $u(1) < 0$ or $u(1) = 0$, $u'(1) > 0$ imply $u'(x_1) < 0$, which means $u(1/x_1) < 0$, which then implies (v) in the same way.

The arguments with y_j's are analogous.

The converse is also straightforward: if (v)(a)–(c) and (vi)(a)–(c) hold, then the above arguments in the reversed order show that $u'(x_j)\overline{u(1/\bar{x}_j)} > 0$ and $u'(y_j)\overline{u(1/\bar{y}_j)} < 0$, which implies that for each such x_j and y_k there exists a *positive* weight w satisfying (5.1). $\square$

6. Point Mass Perturbations

Normally in the theory of orthogonal polynomials we require the measure of orthogonality μ to be of total mass 1. So if we wish to add/remove a pure point to/from the measure, one would have to renormalize the measure by dividing by a constant. We will not do this in the current section just to avoid messiness. Indeed, note that if $p_n(x)$ are orthonormal polynomials for μ, and c is a constant, then $c\mu$ has orthonormal polynomials $p_n(x)/\sqrt{c}$, which satisfy the same recurrence relation as $p_n(x)$. Therefore the associated Jacobi parameters and Jacobi matrix are the same for μ and for $c\mu$.

Until the end of the paper, let μ obey (1.4) or (1.3). Then m has meromorphic continuation to $\mathcal{S}_R$. In particular, m has finitely singularities on $\mathcal{S}_+$. By Lemma 2.7, μ has no singular continuous part. Thus

$$d\mu(x) = f(x)dx + \sum_{j=1}^{N-1} w_j \delta_{z_j}, \tag{6.1}$$

where $\operatorname{ess\,supp}\mu = \operatorname{supp} f(x) = \mathfrak{e}$; $z_j \in \mathbb{R} \setminus \mathfrak{e}$; $w_j > 0$; $1 \le N < \infty$ (we write $N-1$ rather than N just for convenience, since we will be adding an eigenvalue).

Let $(a_n, b_n)_{n=1}^\infty$ be the Jacobi matrix of μ, and $(\widehat{a}_n, \widehat{b}_n)_{n=1}^\infty$ the Jacobi matrix for $\widehat{\mu}$ (where $\widehat{\mu}$ will be defined later). Let m and $\widehat{m}$ be the m-functions of $(a_n, b_n)_{n=1}^\infty$ and $(\widehat{a}_n, \widehat{b}_n)_{n=1}^\infty$, respectively.

6.1. **Removal of point masses.**

Theorem 6.1. *Let μ be* (6.1) *and*

$$d\widehat{\mu}(x) = d\mu(x) - w_{N-1}\delta_{z_{N-1}}.$$

Let $R > 1$. The following holds true.

(i) *If*

$$\limsup_{n\to\infty} \left(|a_n - a_n^{(0)}| + |b_n - b_n^{(0)}| \right)^{1/2n} \le R^{-1},$$

for some periodic Jacobi matrix $(a_n^{(0)}, b_n^{(0)})_{n=1}^\infty$, then

$$\limsup_{n\to\infty} \left(|\widehat{a}_n - \widehat{a}_n^{(0)}| + |\widehat{b}_n - \widehat{b}_n^{(0)}| \right)^{1/2n} \le R^{-1}, \tag{6.2}$$

for some periodic Jacobi matrix $(\widehat{a}_n^{(0)}, \widehat{b}_n^{(0)})_{n=1}^\infty$ from the same isospectral torus.

(ii) *If $(a_n, b_n)_{n=1}^\infty$ is eventually periodic, that is, there exists s such that*

$$a_n = a_n^{(0)}, \quad b_n = b_n^{(0)} \quad \text{for all } n \ge s+1, \tag{6.3}$$

for some periodic Jacobi matrix $(a_n^{(0)}, b_n^{(0)})_{n=1}^\infty$, then

$$\widehat{a}_n = \widehat{a}_n^{(0)}, \quad \widehat{b}_n = \widehat{b}_n^{(0)} \quad \text{for all } n \ge s+1, \tag{6.4}$$

for some periodic Jacobi matrix $(\widehat{a}_n^{(0)}, \widehat{b}_n^{(0)})_{n=1}^\infty$ from the same isospectral torus.

Remark. If $l = 0$ then $(\widehat{a}_n^{(0)}, \widehat{b}_n^{(0)})_{n=1}^\infty = (a_n^{(0)}, b_n^{(0)})_{n=1}^\infty$ of course. Therefore in this case the removal of a mass point is a finite rank perturbation (of rank at most s). If $l \ne 0$ then in general there is no reason for $(\widehat{a}_n^{(0)}, \widehat{b}_n^{(0)})_{n=1}^\infty$ to coincide with $(a_n^{(0)}, b_n^{(0)})_{n=1}^\infty$.

Example. Let us illustrate this on a simple $l = 1$, $s = 0$ case. Suppose $a_{n+2} = a_n$, $b_{n+2} = b_n$ for all n. Instead of writing $(a_n, b_n)_{n=1}^\infty$, let us write simply (a_1, a_2, b_1, b_2). Suppose (a_1, a_2, b_1, b_2) has an eigenvalue $(z_0)_+$ in the gap. If we remove this eigenvalue then the Jacobi matrix $\widehat{\mathcal{J}}$ corresponding to the new measure is going to be the 2-periodic matrix (a_2, a_1, b_1, b_2). Indeed, by [15, Thm 5.2.2], $m^\sharp(z) = (a_2^2 m(z; \mathcal{J}_0^-))^{-1}$, where $\mathcal{J}_0^- = (a_1, a_2, b_2, b_1)$. Thus $m(z; \mathcal{J}_0^-)$ has a zero at $(z_0)_-$. But then once-stripped matrix $(\mathcal{J}_0^-)^{(1)} = (a_2, a_1, b_1, b_2)$ will have a pole at $(z_0)_-$. We know that $\widehat{\mathcal{J}}$ is periodic by the above theorem, and we see that it has the same pole structure as $(\mathcal{J}_0^-)^{(1)} = (a_2, a_1, b_1, b_2)$. This implies that $\widehat{\mathcal{J}} = (a_2, a_1, b_1, b_2)$.

Proof of Theorem 6.1. Note that

$$\widehat{m}(z) = m(z) - \frac{w_{N-1}}{z_{N-1} - z},$$

therefore $\widehat{m}$ has meromorphic continuation to the same domain as m, and

$$\widehat{m}(z) - \widehat{m}^\sharp(z) = m(z) - m^\sharp(z)$$

Thus (ii)(a)–(c) of Lemma 2.5 for $\widehat{m}$ are immediate from (ii)(a)–(c) of Lemma 2.5 for m. Condition (ii)(d) for $\widehat{m}$ is also clear, since the only new pole is at $(z_{N-1})_-$, and $(z_{N-1})_+$ is no longer a pole. $\square$

6.2. **Addition of point masses.** As above, let $\mathcal{J}$ be a Jacobi matrix that is exponentially close to being periodic, (1.4). When adding a point mass at z_N, it is natural to consider two situations: if $(z_N)_-$ is in the domain of meromorphicity of m, and when it is not. The latter case is simpler, and can written as $z_N \in \mathbb{C} \setminus E_R$ in the notation of Lemma 2.5; equivalently,

$$\frac{|\Delta(z_N)| - \sqrt{|\Delta(z_N)|^2 - 4}}{2} \le R^{-1}. \tag{6.5}$$

Note that if $l = 0$ and $\mathfrak{e} = [-2, 2]$ (perturbation of the free Jacobi matrix), then $\Delta(z) = z$, and the left-hand side of (6.5) is the inverse of the map $\mathbb{D} \ni z \mapsto z + z^{-1}$. Let us denote

$$r(z_N)^{-1} = \frac{|\Delta(z_N)| - \sqrt{|\Delta(z_N)|^2 - 4}}{2} \in (0, 1]. \tag{6.6}$$

Theorem 6.2. *Let* μ *be* (6.1) *and*

$$d\widehat{\mu}(x) = d\mu(x) + w_N \delta_{z_N},$$

$z_N \notin \operatorname{supp} \mu$. *Let* $R > 1$ *and*

$$r(z_N)^{-1} \le R^{-1}.$$

If

$$\limsup_{n\to\infty} \left(|a_n - a_n^{(0)}| + |b_n - b_n^{(0)}| \right)^{1/2n} \le R^{-1},$$

for some periodic Jacobi matrix $(a_n^{(0)}, b_n^{(0)})_{n=1}^\infty$, *then*

$$\limsup_{n\to\infty} \left(|\widehat{a}_n - \widehat{a}_n^{(0)}| + |\widehat{b}_n - \widehat{b}_n^{(0)}| \right)^{1/2n} \le R^{-1}, \tag{6.7}$$

for some periodic Jacobi matrix $(\widehat{a}_n^{(0)}, \widehat{b}_n^{(0)})_{n=1}^\infty$ *from the same isospectral torus.*

Theorem 6.3. *Let* μ *be* (6.1) *and*

$$d\widehat{\mu}(x) = d\mu(x) + w_N \delta_{z_N},$$

$z_N \notin \operatorname{supp} \mu$. *Let* $R > 1$ *and*

$$r(z_N)^{-1} > R^{-1}.$$

If

$$\limsup_{n\to\infty} \left(|a_n - a_n^{(0)}| + |b_n - b_n^{(0)}| \right)^{1/2n} \le R^{-1}, \tag{6.8}$$

for some periodic Jacobi matrix $(a_n^{(0)}, b_n^{(0)})_{n=1}^\infty$, *then*

$$\limsup_{n\to\infty} \left(|\widehat{a}_n - \widehat{a}_n^{(0)}| + |\widehat{b}_n - \widehat{b}_n^{(0)}| \right)^{1/2n} \le r(z_N)^{-1} \tag{6.9}$$

for some periodic Jacobi matrix $(\widehat{a}_n^{(0)}, \widehat{b}_n^{(0)})_{n=1}^\infty$ *from the same isospectral torus. Moreover,*

(i) *If* $m(z) = \int \frac{d\mu(x)}{x-z}$ *has a first order pole at* $(z_N)_-$ *and* $\operatorname{Res}_{z=(z_N)_-} m(z) = w_N$, *then*

$$\limsup_{n\to\infty} \left(|\widehat{a}_n - \widehat{a}_n^{(0)}| + |\widehat{b}_n - \widehat{b}_n^{(0)}| \right)^{1/2n} \le R^{-1}. \tag{6.10}$$

(ii) *Otherwise*

$$\limsup_{n\to\infty}\left(|\widehat{a}_n-\widehat{a}_n^{(0)}|+|\widehat{b}_n-\widehat{b}_n^{(0)}|\right)^{1/2n}=r(z_N)^{-1}. \tag{6.11}$$

Remarks. 1. $R=\infty$ is allowed.

2. One can also replace (6.8) and (6.10) with (6.3) and (6.4) respectively. This is of course a different result from just $R=\infty$ case.

3. If $l=0$, then in particular these theorems say that adding or removing a pure point is an exponentially decaying perturbation. If $l\geq 1$, then in general $(\widehat{a}_n^{(0)},\widehat{b}_n^{(0)})_{n=1}^\infty$ is different from $(a_n^{(0)},b_n^{(0)})_{n=1}^\infty$, so the perturbation is not exponentially decaying. One could still say that it is exponentially decaying *with respect to the distance from the isospectral torus.*

4. Note that the (i) of Theorem 6.3 says that adding a pure mass at z_N without modifying the continuous part of μ will work only if z_N is a resonance of order 1, **and** if residue of $m^\sharp$ is positive. From Theorem 3.1, we also now that moving resonance into an eigenvalue has a chance of working only if z_N is an odd-numbered singularity when counted from the nearest band edges. Indeed, by tracking the signs of $m-m^\sharp$, just as in the proof of Theorem 3.1, one can check that positivity of the residue of $m^\sharp$ is equivalent to being an odd-numbered singularity.

Proof of Theorems 6.2 and 6.3. Note that

$$\widehat{m}(z)=m(z)+\frac{w_N}{z_N-z},$$

and therefore $\widehat{m}$ has meromorphic continuation to the same domain as m, and

$$\widehat{m}(z)-\widehat{m}^\sharp(z)=m(z)-m^\sharp(z).$$

Thus (ii)(a)–(c) of Lemma 2.5 for $\widehat{m}$ are immediate from (ii)(a)–(c) of Lemma 2.5 for m.

In the setting of Theorem 6.2, (ii)(d) is also clear, since $(z_N)_-$ is not in $\mathcal{S}_R$ (see (2.1)).

In the setting of Theorem 6.3, condition (ii)(d) for $\widehat{m}$ is holds if and only if $(z_N)_-$ is not a pole of $\widehat{m}(z)$. Thus we need

$$0=\operatorname*{Res}_{z=(z_N)_-}\widehat{m}(z)=\operatorname*{Res}_{z=(z_N)_-}m(z)-w_N.$$

Thus if $w_N=\operatorname{Res}_{z=(z_N)_-}m(z)$ then $\widehat{m}$ satisfies Lemma 2.5(ii) in the same domain as m, so (6.10) holds. Otherwise $\widehat{m}$ satisfies Lemma 2.5(ii) only up to the point $(z_N)_-$. So the maximal r that we apply Lemma 2.5(ii) to, is determined from

$$|\Delta(z_N)|=r+r^{-1},$$

which implies $r^{-1}=r(z_N)^{-1}$ where $r(z)$ is defined in (6.6). □

One can also ask question about what happens when one modifies, rather than removes, the weight at a particular point mass. A very general result of Simon [14, Cor 24.4] says that this always produces an exponentially decaying perturbation:

Lemma 6.4 (Simon)**.** *Let z_0 be an isolated pure point of a measure of compact support μ on $\mathbb{R}$. Let $d\widehat{\mu}(x)=d\mu(x)+\lambda\delta_{z_0}$, where $\lambda>-\mu(\{z_0\})$ so z_0 is also a pure point of $\widehat{\mu}(x)$. Then for some $1>C>0$,*

$$\limsup_{n\to\infty}\left(|a_n-\widehat{a}_n|+|b_n-\widehat{b}_n|\right)^{1/2n}\leq C.$$

Using our methods, assuming that μ satisfies (1.4), we can slightly sharpen the conclusion of this result.

Theorem 6.5. *Suppose that μ satisfies the conditions of Lemma 6.4 and that the associated Jacobi matrix $(a_n, b_n)_{n=1}^\infty$ satisfies*

$$\limsup_{n\to\infty} \Big(|a_n - a_n^{(0)}| + |b_n - b_n^{(0)}|\Big)^{1/2n} = R^{-1}, \tag{6.12}$$

for some periodic Jacobi matrix $(a_n^{(0)}, b_n^{(0)})_{n=1}^\infty$ and some $R > 1$. Then

$$\limsup_{n\to\infty} \Big(|\widehat{a}_n - a_n| + |\widehat{b}_n - b_n|\Big)^{1/2n} = \max\{r(z_0)^{-1}, R^{-1}\}. \tag{6.13}$$

Proof. Indeed, $\widehat{m}$ has singularities at both $(z_0)_+$ and $(z_0)_-$ (unless $\lambda = 0$ or $\lambda = -\mu(\{z_0\})$). Therefore the condition (ii)(d) of Lemma 2.5 breaks down. $\square$

It is amusing that if $\lambda = -\mu(\{z_0\}) + \varepsilon$ with varying $\varepsilon > 0$, then the left-hand side of (6.13) is constant $\max\{r(z_0)^{-1}, R^{-1}\}$. However when $\varepsilon = 0$ then $(\widehat{a}_n, \widehat{b}_n)_{n=1}^\infty$ becomes exponentially close to a completely different periodic matrix (generically), i.e., the left-hand side of (6.13) becomes 1. Of course, even though each Jacobi parameter depends continuously on λ, the $\limsup$ has no reason to be continuous.

7. Christoffel transform of measures

As we have seen in Theorem 6.3, if there is no resonance of m at z_N then we cannot add a pure mass at z_N while preserving the rate of exponential convergence. It was noted by Geronimo (in $l = 0$ situation, see [3, Thm 7]) that if $z_N > \sup\operatorname{supp}\mu$, then one can remedy the situation by first dividing the measure by a linear factor $(z_N - x)$ and then adding the pure mass at z_N. We provide here a simple alternative proof, at the same time generalizing this result to any l.

Theorem 7.1. *Let μ be* (6.1) *and*

$$d\widehat{\mu}(x) = \frac{1}{z_N - x} f(x)dx + \sum_{j=1}^{N-1} \frac{w_j}{z_N - z_j}\delta_{z_j} + w_N\delta_{z_N}, \tag{7.1}$$

where $z_N > \sup\operatorname{supp}\mu$. Let $R > 1$ and

$$\limsup_{n\to\infty} \Big(|a_n - a_n^{(0)}| + |b_n - b_n^{(0)}|\Big)^{1/2n} \le R^{-1}, \tag{7.2}$$

for some periodic Jacobi matrix $(a_n^{(0)}, b_n^{(0)})_{n=1}^\infty$, and

$$w_N = m((z_N)_+) - m((z_N)_-). \tag{7.3}$$

Then

$$\limsup_{n\to\infty} \Big(|\widehat{a}_n - \widehat{a}_n^{(0)}| + |\widehat{b}_n - \widehat{b}_n^{(0)}|\Big)^{1/2n} \le R^{-1} \tag{7.4}$$

for some periodic Jacobi matrix $(\widehat{a}_n^{(0)}, \widehat{b}_n^{(0)})_{n=1}^\infty$ from the same isospectral torus.

Remarks. 1. Again, one can also replace (7.2) and (7.4) with (6.3) and (6.4) respectively.

2. The restriction $z_N > \sup\operatorname{supp}\mu$ is needed to keep the new measure positive. If $z_N < \inf\operatorname{supp}\mu$ then one can similarly divide the measure by $x - z_N$. If z_N does not satisfy either of these conditions then in order to add an eigenvalue at z_N we also need to add a resonance. We discuss this below.

3. Note that there is an implicit restriction here that the right-hand side of (7.3) is positive to ensure $w_N > 0$. In view of Theorem 3.1, we can now understand this restriction: indeed, by tracking the signs of $m - m^\sharp$ in the proof of Theorem 3.1, one sees that $m((z_N)_+) - m((z_N)_-) > 0$ if and only if $[\alpha_{l+1}, z_N)$ (or $(z_N, \beta_l]$) contains an even number of singularities of $\mathcal{J}$! If there is an odd number of them, then one cannot simply add another eigenvalue without the need to add a resonance as well. We address this question later in this section.

Proof. By Lemmas 2.5 and 2.7, $f(x)$ has an meromorphic continuation to $\mathcal{S}_R$. Therefore so does $f(x)/(z_N - x)$. So by Lemma 2.7 again, we obtain that $\widehat{m}$ satisfies Lemma 2.5(ii)(a). (ii)(b) is trivial. Also,

$$m(z) - m^\sharp(z) = 2\pi i f(z), \tag{7.5}$$

$$\widehat{m}(z) - \widehat{m}^\sharp(z) = 2\pi i f(z)/(z_N - x), \tag{7.6}$$

where $f(z)$ is the meromorphic continuation of f to $\pi^{-1}(E_R)$. This shows that if (ii)(c) holds for m, then it holds for $\widehat{m}$.

Let us check (ii)(d) for $\widehat{m}$. Simultaneous poles of $\widehat{m}$ can occur only at $(z_j)_\pm$, where $1 \le j \le N$. If $1 \le j \le N-1$, then by (ii)(d) for m we have $\operatorname{Res}_{(z_j)_+} m^\sharp(z) = 0$, which implies

$$\operatorname*{Res}_{z=(z_j)_+} m(z) = 2\pi i \operatorname*{Res}_{z=(z_j)_+} f(z).$$

Using (7.1), this implies

$$\operatorname*{Res}_{z=(z_j)_+} \widehat{m}(z) = \operatorname*{Res}_{z=(z_j)_+} \frac{2\pi i f(z)}{z_N - x}.$$

Then (7.6) shows that $\widehat{m}^\sharp$ has no pole.

Finally at $(z_N)_+$, the residue of $\widehat{m}$ is $-w_N$, and the residue of $2\pi i f(z)/(z_N - x)$ is $-2\pi i f(z_N) = -\left[m((z_N)_+) - m^\sharp((z_N)_+)\right]$. Since they are equal, cf. (7.3), this means that $m^\sharp$ has no pole at $(z_N)_+$. □

We have already noted (see Remark 4 after Theorem 6.3 and Remark 3 after Theorem 7.1) that it is not always possible to add a mass point to the spectrum without paying attention to the resonances. Indeed, Theorem 3.1 shows that together with adding the eigenvalue, we are usually forced to add a resonance. For example, suppose that a gap (β_1, α_2) contains singularities $z_1 < z_2 < z_3$, and let z_1 and z_3 be eigenvalues and z_2 a resonance. If we want to add an eigenvalue $z_0 \in (z_1, z_2)$, then we also need a resonance on (z_1, z_0) is order for the conditions of Theorem 3.1 to be satisfied. Similarly, if we want to add an eigenvalue $z_0 \in (z_2, z_3)$, then we need a resonance on (z_0, z_3). Moreover, if instead one wants to make z_2 into an eigenvalue, then one needs to add a resonance on (z_1, z_2) and a resonance on (z_2, z_3).

How do we add resonance at a point k_1? This is easily achieved by dividing the measure by $(z - k_1)$. To add an eigenvalue at a point k_2 we divide by $(z - k_2)$ and add $w_2 \delta_{k_2}$ (with w_2 uniquely determined). If k_1 and k_2 are close together this preserves the positivity of the measure.

Finally, suppose we want to add an eigenvalue at k_1 while there is a resonance of order $N \ge 1$ at k_1. Then we should remove the resonance (by multiplying the measure by $(z - k_1)^N$). To keep the measure positive (equivalently, to satisfy the condition (iv) of Theorem 3.1), we should immediately introduce new N (or N

mod 2 is enough) resonances close to k_1 by dividing the measure by $(z-(k_1\pm\varepsilon))^N$. Now k_1 is no longer a singularity and so we can apply the above procedure.

We formalize these ideas in the next theorem.

Theorem 7.2. *Let μ be (6.1), $R>1$, and the associated Jacobi matrix satisfies*

$$\limsup_{n\to\infty}\Big(|a_n-a_n^{(0)}|+|b_n-b_n^{(0)}|\Big)^{1/2n}\le R^{-1} \tag{7.7}$$

for some periodic Jacobi matrix $(a_n^{(0)},b_n^{(0)})_{n=1}^\infty$.

Let $z_N\notin\operatorname{supp}\mu$. Then there exists a rational function $g(x)$ and $w_N>0$ such that measure

$$d\widehat{\mu}(x)=g(x)f(x)dx+\sum_{j=1}^{N-1}g(z_j)w_j\delta_{z_j}+w_N\delta_{z_N}, \tag{7.8}$$

is positive, and satisfies

$$\limsup_{n\to\infty}\Big(|\widehat{a}_n-\widehat{a}_n^{(0)}|+|\widehat{b}_n-\widehat{b}_n^{(0)}|\Big)^{1/2n}\le R^{-1} \tag{7.9}$$

for some periodic Jacobi matrix $(\widehat{a}_n^{(0)},\widehat{b}_n^{(0)})_{n=1}^\infty$ from the same isospectral torus.

Proof. Suppose first that there is no resonance of μ at z_N. Choose $\varepsilon>0$ small enough so that there is no singularity of μ in $(z_N-\varepsilon,z_N+\varepsilon)$. Then let

$$g(x)=\frac{1}{(x-z_N)(x-z_N\pm\varepsilon)},$$

where the sign is to be determined later. It is clear from the proof of Theorem 7.1 that the measure

$$d\nu(x)=g(x)f(x)dx+\sum_{j=1}^{N-1}g(z_j)w_j\delta_{z_j} \tag{7.10}$$

satisfies conditions (ii)(a)–(d) of Lemma 2.5, and contains the same system of singularities as μ plus two new first order resonances at z_N and $z_N\mp\varepsilon$. Let $n(z)$ be the m-function of ν. Now let us apply Theorem 6.3(i) to ν at z_N. It can be applied only if

$$w_N=\operatorname*{Res}_{z=(z_N)_+}n^\sharp(z)$$

is positive. But by Lemma 2.7 and the fact that $(z_N)_+$ is not an eigenvalue, we get

$$\operatorname*{Res}_{z=(z_N)_+}n^\sharp(z)=-2\pi i\operatorname*{Res}_{z=(z_N)_+}\frac{f(z)}{(z-z_N)(z-z_N\pm\varepsilon)}=\frac{-2\pi i f((z_N)_+)}{\pm\varepsilon}.$$

Note that $f((z_N)_+)$ is non-zero by Lemma 2.5(ii)(c). Also, $if((z_N)_+)$ is real since both m and $m^\sharp$ are real on $\pi^{-1}(\mathbb{R}\backslash\mathfrak{e})$. Therefore choosing either positive or negative sign in front of ε, we can make sure that $w_N>0$.

Now suppose that z_N is a resonance of of μ order $k\ge1$. Choose $\varepsilon>0$ small enough so that there is no singularity of μ in $(z_N-\varepsilon,z_N+\varepsilon)$ other than z_N. Then let

$$g(x)=\frac{(x-z_N)^{k-1}}{(x-z_N-\varepsilon)^{k_1}(x-z_N+\varepsilon)^{k+1-k_1}}.$$

Note that g is positive on $\operatorname{supp}\mu$. The same arguments as before show that the measure

$$d\nu(x) = g(x)f(x)dx + \sum_{j=1}^{N-1} g(z_j)w_j\delta_{z_j} \tag{7.11}$$

satisfies conditions (ii)(a)–(d) of Lemma 2.5, and contains the same system of singularities as μ except for the new resonances of order k_1 at $z_N+\varepsilon$, of order $k+1-k_1$ at $z_N-\varepsilon$, and the order of the resonance at z_N is now 1 instead of k. Now we can apply Theorem 6.3(i) to ν at z_N, and see that

$$\begin{aligned} w_N = \operatorname*{Res}_{z=(z_N)_+} n^\sharp(z) &= -2\pi i \lim_{z\to(z_N)_+}(z-z_N)\frac{(z-z_N)^{k-1}f(z)}{(z-z_N-\varepsilon)^{k_1}(z-z_N+\varepsilon)^{k+1-k_1}} \\ &= (-1)^{k_1-1}\frac{2\pi i \lim_{z\to(z_N)_+}(z-z_N)^k f(z)}{\varepsilon^{k+1}} \end{aligned}$$

Now $i\lim_{z\to(z_N)_+}(z-z_N)^k f(z)$ is real, and so choosing either $k_1=0$ or 1 will make $w_N>0$. □

Using the same approach it is easy to see that we can also modify resonances in any way we want if it does not violate the conditions of Theorem 3.1.

Corollary 7.3. *Let $R>1$, and the associated to a measure μ Jacobi matrix satisfies*

$$\limsup_{n\to\infty}\left(|a_n-a_n^{(0)}|+|b_n-b_n^{(0)}|\right)^{1/2n} \le R^{-1} \tag{7.12}$$

for some periodic Jacobi matrix $(a_n^{(0)},b_n^{(0)})_{n=1}^\infty$.

For any system of singularities that is consistent with Theorem 3.1 and that differs from the system of singularities of μ in finitely many points, there exists a Christoffel transform and a sequence of Uvarov transforms that produces a positive measure $\widehat{\mu}$ with such a system of singularities.

Remarks. 1. To destroy a resonance or an eigenvalue at z_0 one needs to multiply the measure by $(z-z_0)$. To keep the measure positive (equivalently, to keep Theorem 3.1 satisfied), one also needs to delete a neighboring singularity, or to introduce another one.

2. By an Uvarov transform we mean addition of a new point mass, and by a Christoffel transform, we mean (7.11).

8. Perturbations of block Jacobi matrices

Theorems 6.1, 6.2, 6.3, 7.1 for $l=0$ can be generalized to the matrix-valued setting. By this we mean the (block) Jacobi matrices with the coefficients a_n and b_n being $m\times m$ matrices. The approach to the proofs is the same, except instead of Lemmas 2.5 and 2.6 we need to use their matrix-valued extensions. These are proved by the author in [12, Thm 3.8–3.9]. The main difference is that (ii)(d) gets more complicated because now simultaneous poles at z and $z^\sharp$ are possible but under additional constraints.

The statement of these matrix-valued generalization for Theorems 6.1, 6.2, 7.1 is exactly identical to the scalar versions, except now $\mathfrak{e}=[-2,2]$, $\Delta(z)=z$, instead of (1.4) and (1.3) one now has

$$\limsup_{n\to\infty}\left(||b_n||+||\mathbf{1}-a_na_n^*||\right)^{1/2n} \le R$$

and

$$b_n = \mathbf{0}, \quad a_n a_n^* = \mathbf{1} \quad \text{for all } n \geq s+1,$$

respectively. Here $\mathbf{1}$ and $\mathbf{0}$ are $m \times m$ identity and zero matrices, and a^* is the Hermitian conjugate.

The matrix-valued analogue of Theorem 6.3 slightly differs from scalar case:

Theorem 8.1. *Let μ be a (matrix-valued) measure* (6.1) *and*

$$d\widehat{\mu}(x) = d\mu(x) + w_N \delta_{z_N},$$

$z_N \notin \operatorname{ess\,supp} \mu = [-2,2]$. *Let $R > 1$ and*

$$r(z_N)^{-1} > R^{-1}.$$

If

$$\limsup_{n\to\infty} \left(||b_n|| + ||\mathbf{1} - a_n a_n^*||\right)^{1/2n} \leq R \tag{8.1}$$

then

$$\limsup_{n\to\infty} \left(||b_n|| + ||\mathbf{1} - a_n a_n^*||\right)^{1/2n} \leq r(z_N)^{-1} \tag{8.2}$$

Moreover,

(i) *If m has no singularity at $(z_N)_-$, then*

$$\limsup_{n\to\infty} \left(||\widehat{b}_n|| + ||\mathbf{1} - \widehat{a}_n \widehat{a}_n^*||\right)^{1/2n} = r(z_N)^{-1} \tag{8.3}$$

(ii) *If $m(z) = \int \frac{d\mu(x)}{x-z}$ has a first order pole at $(z_N)_-$, then*

$$\limsup_{n\to\infty} \left(||\widehat{b}_n|| + ||\mathbf{1} - \widehat{a}_n \widehat{a}_n^*||\right)^{1/2n} = \limsup_{n\to\infty} \left(||\widehat{b}_n|| + ||\mathbf{1} - \widehat{a}_n \widehat{a}_n^*||\right)^{1/2n} \leq R \tag{8.4}$$

if and only if $w_N = P\left(\operatorname{Res}_{z=(z_N)_-} m(z)\right) P$, where P is the orthogonal projection onto an invariant subspace of $\operatorname{Res}_{z=(z_N)_-} m(z)$. If w_N is not of this form then (8.3) *holds.*

(iii) *If $m(z) = \int \frac{d\mu(x)}{x-z}$ has a pole of order higher than 1 at $(z_N)_-$, then* (8.4) *holds if and only if*

$$\operatorname{Ran} w_N \subset \ker\left[(m((z_N)_-) - m^\sharp((z_N)_-))^{-1}\right],$$

$$\operatorname{Ran} w_N \subset \left(\operatorname{Ran} \lim_{z\to(z_N)_-} \left((m(z) - m^\sharp(z))^{-1} m(z) + (m(z) - m^\sharp(z))^{-1} \frac{w_N}{z - z_N}\right)\right)^\perp$$

Otherwise (8.3) *holds.*

The proofs of these results are written in [11, Sect 1.3].

Acknowledgement. The author would like to thank Rowan Killip for useful discussions.

References

[1] David Damanik and Barry Simon. Jost functions and Jost solutions for Jacobi matrices. II. Decay and analyticity. *Int. Math. Res. Not.*, Art. ID 19396, 32 pages, 2006.

[2] Iryna Egorova, Johanna Michor, and Gerald Teschl. Scattering theory for Jacobi operators with quasi-periodic background. *Comm. Math. Phys.*, 264(3):811–842, 2006.

[3] Jeffrey S. Geronimo. Scattering theory, orthogonal polynomials, and q-series. *SIAM J. Math. Anal.*, 25(2):392–419, 1994.

[4] Jeffrey S. Geronimo and Paul G. Nevai. Necessary and sufficient conditions relating the coefficients in the recurrence formula to the spectral function for orthogonal polynomials. *SIAM J. Math. Anal.*, 14(3):622–637, 1983.

[5] Fritz Gesztesy and Gerald Teschl. Commutation methods for Jacobi operators. *J. Differential Equations*, 128(1):252–299, 1996.

[6] David S. Greenstein. On the analytic continuation of functions which map the upper half plane into itself. *J. Math. Anal. Appl.*, 1:355–362, 1960.

[7] Alexei Iantchenko and Evgeny Korotyaev. Periodic Jacobi operator with finitely supported perturbation on the half-lattice. *Inverse Problems*, 27(11):115003, 26, 2011.

[8] Rowan Killip and Barry Simon. Sum rules for Jacobi matrices and their applications to spectral theory. *Ann. of Math. (2)*, 158(1):253–321, 2003.

[9] E. L. Korotyaev. Inverse resonance scattering for Jacobi operators. *Russ. J. Math. Phys.*, 18(4):427–439, 2011.

[10] Rostyslav Kozhan. Meromorphic continuations of finite gap Herglotz functions and periodic orthogonal polynomials. (under submission), arXiv:1210.4627).

[11] Rostyslav Kozhan. *Asymptotics for orthogonal polynomials, exponentially small perturbations and meromorphic continuations of Herglotz functions*. Ph.D. thesis, California Institute of Technology, Pasadena, CA, 2010.

[12] Rostyslav Kozhan. Jost asymptotics for matrix orthogonal polynomials on the real line. *Constr. Approx.*, 36(2):267–309, 2012.

[13] Barry Simon. Resonances in one dimension and Fredholm determinants. *J. Funct. Anal.*, 178(2):396–420, 2000.

[14] Barry Simon. The Christoffel-Darboux kernel. In *Perspectives in partial differential equations, harmonic analysis and applications*, volume 79 of *Proc. Sympos. Pure Math.*, pages 295–335. Amer. Math. Soc., Providence, RI, 2008.

[15] Barry Simon. *Szegő's theorem and its descendants: spectral theory for L^2 perturbations of orthogonal polynomials*. M. B. Porter Lectures. Princeton University Press, Princeton, NJ, 2011.

[16] Manwah Lilian Wong. Point mass insertion on the real line and non-exponential perturbation of the recursion coefficients. *Math. Nachr.*, 285(2-3):377–388, 2012.

[17] Maciej Zworski. Distribution of poles for scattering on the real line. *J. Funct. Anal.*, 73(2):277–296, 1987.

E-mail address: `kozhan@math.ucla.edu`

University of California, Los Angeles, Department of Mathematics, Los Angeles, CA 90095, USA